THE VENUS FLAGSHIP STUDY: ATMOSPHERIC SCIENCE FROM TWO DESCENT/LANDING VEHICLES AND TWO BALLOONS

Mark A. Bullock¹, Jeffrey L. Hall² and Tibor S. Balint²

¹Southwest Research Institute ²Jet Propulsion Laboratory





NASA'S Flagship Mission to Venus April 6, 2009





One year study

Report released 6 April 2009

Team consisted of atmospheric scientists, planetary geologists, planetary geophysicists, mission designers, technology experts, and engineers

Important architecture and costing support by JPL's Team X

Beautiful web site!


http://vfm.jpl.nasa.gov/

Venus STDT Objectives

- Phase 1: Develop and Prioritize Science Objectives and Investigations for a Venus flagship mission drawing upon
 - VEXAG White Paper (2007)
 - NRC Decadal Survey (2003) and NOSSE update (2008)
 - NASA's SSE Roadmap (2006)
- Phase 1: Identify the optimal architecture to achieve science objectives
- Phase 2: Execute the Design Reference Mission design from a science-driven architecture trade
 - Design and assess scientific performance of payload
 - Assess Performance, Cost, Risk, and Technology Readiness
- Phase 2: Identify Technology Development
 - To fully execute the Design Reference Mission
 - For alternate payloads and architectures

International Collaboration

- Multi-element architecture lends itself to international collaboration
- Timing for international collaboration:
 - NASA (Venus Flagship)
 - ESA (VEX Current-2011)
 - ESA (Cosmic Vision EVE > 2020)
 - JAXA (VCO 2010)
 - Russia (Venera D 2016)



Venus Flagship Mission Assumptions & Constraints

Launch Opportunity: 2020 to 2025

Technology Maturation: TRL 6 by 2015

Life Cycle Mission Cost Range: \$3 - 4B (FY '08)

LV Capability: ≤ Delta IVH equiv.

DSN Capability: up to 34M, Ka band

International Contribution:
 No foreign cost contribution


 Assume no earlier missions prior to flight of the Venus Flagship Mission

Why is Venus so different from Earth?


- What does the Venus greenhouse tell us about climate change?
 - How do clouds and chemical cycles affect atmospheric energy balance?
 - What drives the atmospheric superrotation?
 - Is there evidence for climate change?
- How active is Venus?
 - Is Venus geologically active and what is it's geologic history?
 - How do surface/atmosphere interactions affect rock mineralogy and climate?
 - What is structure of the interior, and what are its dynamics?
- When and where did the water go?
 - How did the early atmosphere evolve?
 - Did Venus have an ocean and if so, when was it lost?
 - Is there continent-like crust on Venus?


The Venus Greenhouse

How does the greenhouse work, and has Venus experienced climate changes?




Mariner 10






Is Venus Geologically Active?

Plains



How did early Venus evolve? Did it have oceans?




High D/H means
Venus once had much
more water.


Venus' climate is an interconnected system of atmospheric, geologic, and surface chemistry processes, just like the Earth's.

Was Venus once habitable?

Learning about Earth by Studying Venus

Prediction of O₃ loss due to CFCs followed directly from studying Venus atmospheric chemistry. TOMS below (Earth)





South polar vortex of Venus exhibits instability and ways to test Earth atmosphere dynamics VEX (left IR emission, right solar reflection)

Venus Flagship Science Objectives

Science Theme	Science Objective
	Understand radiation balance in the atmosphere and the cloud and
What does the Venus greenhouse	chemical cycles that affect it
tell us about climate change?	Understand how superrotation and the general circulation work
	Look for evidence of climate change at the surface
	Identify evidence of current geologic activity and understand the
How active is Venus?	geologic history
	Understand how surface/atmosphere interactions affect rock
	chemistry and climate
	Place constraints on the structure and dynamics of the interior
	Determine how the early atmosphere evolved
When and where did the water go?	Identify chemical and isotopic signs of a past ocean
where did the water go:	Understand crustal composition differences and look for evidence of
	continent-like crust


Main resources:

- •2003 Decadal Survey
- •VEXAG 2007 report
- •2006 NASA Roadmap

Mission Architecture Trade Study


- STDT & JPL Team assessed the mission architecture trade space:
 - Identified 13 architectural elements
 - e.g. orbiter, landers, balloons, probes
 - Targeting various altitude regimes
 - from surface to low/mid/high altitudes and orbit
 - Single and multiple elements were considered
 - e.g., networks, multi-probes
- These architecture elements assembled into 17 mission architectures
- Science FOMs and Technology Difficulty used to select highest value science architecture within design constraints

Magellan - Ishtar Terra




Design Reference Mission

- The DRM requires a dual-launch approach using a pair of Atlas V 551s
 - 1 orbiter, arrives first to serve as telecom relay
 - 1 carrier, arrives second, with 2 entry vehicles, each with a balloon and a lander, delivered into the atmosphere 13 hours apart
 - Launches in 2021, arrivals in 2022
 - Orbiter serves as a telecom relay for landers (5 hours) and balloons (30 days)
 - 2 year radar mapping science mission after aerobraking to a 230 km circular orbit
- The balloons and landers communicate through the orbiter, with the carrier serving as a limited emergency backup
- The lander sites are Alpha Regio (-27°, 3°
 E) and the lava flows at -47.4°, 6.5°
 - The balloons are expected to circumnavigate Venus 5-7 times and drift poleward








DRM Planning Payload


Orbiter	2 Balloons	2 Landers					
Lifetime (4 years)	(1 month)	Descent Phase (1–1.5 hour)	Landed Phase (5 hours)				
InSAR — Interferometric Synthetic Aperture Radar	ASI — Atmospheric Science Instrument (pressure, temperature, wind speed,)	ASI	Microscopic imager				
Vis-NIR Imaging Spectrometer	GC/MS — Gas Chromatograph / Mass Spectrometer	Vis-NIR Cameras with spot spectrometry	XRD / XRF				
Neutral Ion Mass Spectrometer	Nephelometer	GC / MS	Heat Flux Plate				
Sub-mm Sounder	Vis-NIR camera	Magnetometer	Passive Gamma Ray Detector				
Magnetometer	Magnetometer	Net Flux Radiometer	Sample acquisition, transfer, and preparation				
Langmuir Probe	Radio tracking	Nephelometer	Drill to ~10 cm				
Radio Subsystem (USO — Ultra Stable Oscillator)			Microwave corner reflector				

Mission Design




- Carrier with entry vehicles launches first, arrives second
- Orbiter launches second, arrives first and sets up to be a telecom relay in a 300 x 40000 km near polar orbit
- Entry vehicles arrive 13
 hours apart so only 1 lander
 communicates with the
 orbiter at a time
- After 1 month of balloon mission, orbiter ends telecom support and aerobrakes down to a 230 km circular orbit for a 2 year science mission phase

Carrier and Entry Vehicles




The carrier uses an inline stacked configuration for the entry vehicles. There is ample room inside the 4.5 m launch vehicle fairing.

The entry vehicles are PV and Galileo style 45° sphere cone aeroshells, sized at 2.65 m diameter to accommodate all internal components.



Landers and Balloons








5.5 m prototype superpressure balloon (JPL/ILC Dover/NASA Wallops)

Entry, Descent and Landing (Inflation)



Launch Mass Summary

• Each of the two launch vehicles in our architecture needed to be the largest Atlas V available (Atlas 551):

• Orbiter:

■ Dry Mass (CBE) = 1591 kg

■ Margin (43%) = 684 kg

■ Propellant = 3030 kg

■ Total = 5305 kg

■ LV capability = 5450 kg

■ LV Margin = 2.6%



■ Carrier (CBE) = 781 kg

Entry Vehicles (2) CBE = 1566 kg

■ Landers (2) CBE = 962 kg

■ Balloons (2) CBE = 209 kg

■ Helium CBE = 26 kg


■ Margin (43%) = 1511 kg

■ Propellant = 523 kg

■ Total = 5578 kg

■ LV capability = 5580 kg

■ LV Margin = 0%



Data Volume Summary

- Preliminary data collection budgets were developed for all of the instruments on all of the platforms.
 - Orbiter (300 Tbits):
 - InSAR instrument provides ~99.99% of all orbiter data
 - Lander (1 Gbit each):

Panoramic imaging: 590 Mbits (59%)
Descent imaging: 200 Mbits (20%)
XRD/XRF: 140 Mbits (14%)

Balloon (21 Mbit each):

VASI + nephelometer: 18 Mbits (86%)
GCMS: 1.6 Mbits (8%)
Microphone 0.7 Mbits (3%)

- The data collection strategy is different for each platform:
 - Orbiter data is collected throughout the 2 year main science mission
 - Lander data is collected continuously during the 1 hour descent and 5 hour surface mission.
 - Balloon data is collected over 30 days, with significant duty-cycling to conserve electrical power.

Primary Issues and Risks

- Sample acquisition and handling
 - Venera/VEGA heritage is dated, improved capabilities likely will be required.
- Lander design and technologies
 - Require design for safe landing on rough terrain (tessera). Rotating pressure vessel concept requires development and validation.
- Launch vehicle limits
 - Atlas 551 limit already reached with 43% margin on CBE. Further mass growth will require descopes or much more expensive launch vehicles.
- Orbiter failure risk
 - Carrier can provide only a limited backup telecom capability if the orbiter fails.
- System engineering complexity
 - The multi-element architecture is complex and few system engineering details have been worked out so far.
- Cost estimation uncertainties
 - Lack of experience and existing facilities makes it difficult to estimate costs for high temperature and high pressure V&V of lander and exposed instruments.

DRM Cost Estimates

- The JPL cost-complexity model (Peterson et al, 2008) gave a \$2.7B estimate for the Design Reference Mission
 - The stated accuracy of this model ~40% on absolute cost, implying a maximum possible cost of \$3.8B.
- The JPL study team created a second cost estimate that fused three sources: JPL's Team X cost models, study team expert inputs and a cost risk subfactor analysis to determined a recommended reserve level. The result is:

Element	Cost
	(\$M)
Spacecraft CBE from Team X	1954
Additional PSE costs	30
Cost Reserves on CBE (41%)	813
Two Atlas 551 launch vehicles	445
Additional Technology Development Costs	107
Total	3349

- This cost is \$3.35B, which is within the uncertainty range of the cost-complexity model.
- The final report lists a cost range of \$2.7B to \$3.8B for the DRM

Technology Priorities

	Technologies for DRM
1	Surface sample acquisition system and handling at Venus surface
2	Lander technologies for rotating pressure vessel and rugged terrain survivability
3	Venus-like environmental test chamber
	New capabilities
4	Refrigeration for the Venus surface environment
5	High temperature sensors and electronics, including telecom systems
	Enhancement to current DRM design
6	Extension of lander life through advanced thermal control

DRM Science Traceability- Themes to Observation Platforms

Science Theme	Science Objective	Instrument Type	Observation Platform		
	Understand radiation balance in the atmosphere	Vis-NIR Imaging Spectrometer	Orbiter		
	and the cloud and chemical cycles that affect it	Nephelometer	Balloon plus Lander (on descent)		
		Net Flux Radiometer	Lander (on descent)		
What does the Venus greenhouse	Understand how superrotation and the general	Sub-millimeter Sounder	Orbiter		
tell us about climate change?	circulation work	Atmospheric Structure (P/T/winds/accel)	Balloon plus Lander (on descent)		
		Radio (with USO)	Baloon		
	Look for evidence of climate change at the surface	Vis-NIR Camera	Balloon plus Lander (on descent)		
		Microscopic Imager	Lander		
	Identify evidence of current geologic activity and	InSAR	Orbiter		
	understand the geologic history				
	Understand how surface/atmosphere interactions	GC/MS	Lander (on descent)		
How active is Venus?	affect rock chemistry and climate				
now active is verius?	Place constraints on the structure and dynamics of	Radio (with USO)	Orbiter		
	the interior	Magnetometer	Orbiter, Balloon, Lander		
		Heat Flux Plate	Lander		
		Corner Reflector	Lander		
	Determine how the early atmosphere evolved	GC/MS	Balloon plus Lander (on descent)		
		Langmuir Probe	Orbiter		
		Neutral and Ion Mass Spectrometer (INMS)	Orbiter		
When and where did the water go?	Identify chemical and isotopic signs of a past ocean		Lander		
		Drill and sample acquisition, transfer and	Lander		
		preparation			
	·	Passive Gamma-ray Detector	Lander		
	look for evidence of continent-like crust				

Open Science Questions

The Atmosphere

MAJOR OPEN SCIENTIFIC QUESTIONS ABOUT VENUS	DESIGN REFERENCE MISSION			ON
	Orbiter	Landers	Balloons	DRM with synergies
VENUS ATMOSPHERE				
How did Venus evolve to become so different from Earth?				
Was Venus ever habitable, and for how long? Did Venus lose a primary atmosphere due to impacts or loss to space?				
What drives Venus' atmospheric superrotation?				
How do geologic activity and chemical cycles affect the clouds and climate?				
How are atmospheric gases lost to space?				
	Fully a	ddresses ques	stion	
Major progress in answering the of the questions Will not answer the questions			swering the q	uestion
			questions	

Open Science Questions Geology

MAJOR OPEN SCIENTIFIC QUESTIONS ABOUT VENUS		DESIGN REFERENCE MISSION			ON
		Orbiter	Landers	Balloons	DRM with synergies
VENUS GEOLOGY					
What is the volcanic and tectonic resurfacing history of Venus?					
How were the heavily deformed highlands made?					
How active is Venus geologically?					
Did Venus ever have plate tectonics and if so, when did it cease?				_	
How are geology and climate connected on Venus?					
What has been the role of water and other volatiles in Venus geology?					
		Fully add	dresses ques	tion	
Major progress in answering the			swering the q	uestion	
	Partial answer to the questions				
	Will not answer the questions				

Open Science Questions

Interior Structure

MAJOR OPEN SCIENTIFIC QUESTIONS ABOUT VENUS		DESIGN REFERENCE MISSION			ION
	Or	biter	Landers	Balloons	DRM with synergies
VENUS INTERIOR STRUCTURE Does Venus have Earth-like continents? What are the chemical, physical, and thermal conditions of the interior? How does mantle convection work on Venus? What is the size and physical state of the core? What is the structure of the Venus lithosphere? How have water and other volatiles affected Venus' interior evolution?					
	N Pa	Fully addresses question Major progress in answering the question Partial answer to the questions Will not answer the questions			

Open Science Questions Geochemistry


MAJOR OPEN SCIENTIFIC QUESTIONS ABOUT VENUS		DESIGN REFERENCE MISSION			ION
	0	rbiter	Landers	Balloons	DRM with synergies
VENUS GEOCHEMISTRY Was there ever an ocean on Venus, and if so, when and how did it disappear? What caused the resufacing of Venus over the past billion years? What is the nature of chemical interactions between surface and atmosphere? What are the tectonic forces behind Venus' volcanism? How were the rocks and soils of Venus formed? What do chemical differences of terrains say about the evolution of Venus?					
	N F	Major pr Partial a	dresses ques ogress in ans nswer to the answer the	swering the c	juestion



- Elemental & mineralogical analysis of rocks & soils
- Descent & panoramic imaging
- Deep atmosphere composition
- Corner reflector, heat flux


Venus Flagship In-Situ Science

- Winds, cloud chemistry and microphysics
- Noble gases
- Radiative energy balance



STDT Summary

- What does the Venus greenhouse tell us about climate change?
 - Probes through atmosphere simultaneously with balloons
 - Chemistry of the surface
- How active is Venus?
 - Highly capable orbiter with high resolution radar imaging, topography, and temporal changes
 - Geothermal heat flux
 - Near-IR images
- When and where did the water go?
 - Geochemistry and mineralogy at 2 locations on Venus
 - Atmospheric isotopes for early evolution
- A Venus Flagship mission in 2020-2025 can be done with moderate technology investment and relatively low risk.



Conclusions

 An orbiter, 2 balloons and 2 landers provide the highest science return.



- VFM studies Venus as an interconnected system atmosphere, clouds, surface, and interior.
- VFM is a large NASA planetary mission with the explicit intention of better understanding our own world.





Venus STDT Membership

Chairs: Mark Bullock (SwRI) and Dave Senske (JPL)

• Atmosphere

- David Grinspoon (DMNS)
- Anthony Colaprete (NASA Ames)
- Sanjay Limaye (U. Wisconsin)
- George Hashimoto (Japan)
- Dimitri Titov (ESA)
- Eric Chassefiere (France)
- Hakan Svedhem (ESA)

Geochemistry

- Allan Treiman (LPI)
- Steve Mackwell (LPI)
- Natasha Johnson (NASA)

Geology and Geophysics

- Dave Senske (JPL)
- Jim Head (Brown University)
- Bruce Campbell (Smithsonian)
- Gerald Schubert (UCLA)
- Walter Kiefer (LPI)
- Lori Glaze (GSFC)

Technology

- Elizabeth Kolawa (JPL)
- Viktor Kerzhanovich (JPL)
- Gary Hunter (GRC)
- Steve Gorevan (HoneyBee)

Ex Officio

- Ellen Stofan (VEXAG Chair)
- Tibor Kremic (NASA)

JPL Venus Flagship Mission Architecture Study

Study Lead: Jeff Hall

Tibor Balint

Craig Peterson

Alexis Benz

Team X

NASA and JPL

Jim Cutts (JPL)

Adriana Ocampo (NASA HQ)